

NUTRITIONAL STATUS AND IMMUNE FUNCTION  
IN COCAINE AND HEROIN ABUSERS AND  
IN METHADONE TREATED SUBJECTS

N. D. Huggins,<sup>1</sup> M. A. Khaled,<sup>1,2</sup> P. E. Cornwell<sup>2</sup>  
and J. O. Alvarez,<sup>2,3</sup>

Departments of <sup>1</sup>Psychiatry and Behavioral Neurobiology,  
<sup>2</sup> Nutrition Sciences, and <sup>3</sup>Public Health Sciences,  
University of Alabama at Birmingham, University Station,  
Birmingham, AL 35294, U.S.A

ABSTRACT

The plasma levels of some essential nutrients and the lymphocyte CD<sub>4</sub> to CD<sub>8</sub> ratio were measured in four groups of individuals that included: (a) cocaine and (b) heroin abusers, (c) methadone treated and (d) healthy subjects. Folate and B-carotene levels were lower in the three drug groups while vitamin C was lower in the methadone and heroin subjects. Vitamin E levels were borderline low in the methadone and cocaine groups. The methadone group also showed a significantly higher level of lipid peroxidation which correlated well with the low values observed for the antioxidant nutrients. Interestingly, the methadone group was the only one with a significantly reduced lymphocyte CD<sub>4</sub>/CD<sub>8</sub> ratio.

INTRODUCTION

Drug users, particularly intravenous drug abusers, are at significant risk of contracting HIV, hepatitis B and other infections. Chandra, 1983 and numerous other authors have shown

that malnutrition and infection are mutually aggravating through their common association with immunity. Also, malnutrition has been linked to the reduction of lymphocyte T-helper (CD<sub>4</sub>) to T-suppressor (CD<sub>8</sub>) ratio (Casey *et al.*, 1983). On the other hand, addictive drugs of abuse have been shown not only to produce significant nutritional aberrations (Mohs, Watson and Leonard-Green, 1990), but can also modulate immunocompetence (Donahoe and Falek, 1988).

Almost 25 years ago, methadone maintenance programs were started to prevent drug abusers from taking hard-core narcotic drugs. Methadone, however, is a psychoactive drug, which ranks 17th on the Drug Abuse Warning Network, albeit far below heroin. Immunological abnormalities have been reported to occur frequently in heroin addicts and in methadone treated subjects (Kreek, 1990) although Novick *et al.*, 1989 have reported that subjects on methadone programs for more than 11 years have normal immune functions in terms of CD<sub>4</sub>/CD<sub>8</sub> T-cell ratio. Recently, however, Herbert *et al.*, 1990 reported nutritional abnormalities in terms of body mass index (BMI), hematocrit and hemoglobin concentration in patients on a methadone maintenance program.

Nutritional status in general and specific nutrient imbalances in particular have been shown to affect the immune response. It is plausible that nutritional deficits, common in drug abusers, may mediate or contribute to the deterioration of the immune function in these individuals. Since there is a lack of information in the literature, a preliminary study was designed in order to gain some insight into this problem.

METHODS

Twenty-four human subjects participated in this study to compare four groups of six subjects: group 1, healthy subjects as control; group 2, methadone treated for more than six months and less than five years; group 3, heroin addicts prior to entering into the methadone maintenance program; and group 4, cocaine abusers. None of the subjects were HIV positive and the human use protocol was approved by the Institutional Review Board of the University of Alabama at Birmingham (UAB). Blood samples were withdrawn from each subject for the analysis of some essential micronutrients and to estimate their immune status in terms of CD<sub>4</sub>/CD<sub>8</sub> including lipid peroxidation (as MDA) and glutathione (GSH) levels.

Vitamin levels were determined using standard laboratory procedures in UAB's Nutrition Diagnostic Laboratory while the measurements of CD<sub>4</sub> and CD<sub>8</sub> cells were made by the fluorescence technique (Fahey, et al., 1984) using FITC conjugated OKT4 and OKT8, respectively. The GSH level was measured spectrophotometrically, (Beutler et al., 1963) and the lipid peroxidation level in terms of malondialdehyde (MDA) production was estimated by HPLC (Sato, 1978). Statistical calculations were performed by using ANOVA analysis and the significant differences are reported in the table.

TABLE

	Group 1 <u>Control</u>	Group 2 <u>Methadone</u>	Group 3 <u>Oplate</u>	Group 4 <u>Cocaine</u>
Plasma Folate (ng/ml)	13.98 ± 6.85	6.73 ± 3.79*	4.48 ± 1.87**	6.16 ± 6.53
RBC Folate (ng/ml)	653.17 ± 218.46	378.83 ± 88.07*	336.33 ± 53.34**	217.17 ± 111.23**
Vitamin B12 (pg/ml)	385 ± 108.9	930.67 ± 577.86	447.67 ± 45.74	577.5 ± 289.60
Vitamin B6 (A/C)	1.46 ± 0.18	1.70 ± 0.20	1.76 ± 0.26*	1.76 ± 0.34
Riboflavin (A/C)	1.22 ± 0.11	1.19 ± 0.07	1.37 ± 0.21	1.56 ± 0.29*
Thiamine (A/C)	1.08 ± 0.06	1.13 ± 0.04	1.13 ± 0.03	1.11 ± 0.05
Vitamin C (mg/dl)	0.76 ± 0.26	0.20 ± 0.14**	0.05 ± 0.07**	0.44 ± 0.44
Vitamin A (ug/dl)	51.5 ± 13.4	42.58 ± 13.79	53.67 ± 17.34	51.33 ± 19.77
Carotenes (ug/dl)	170.17 ± 53.18	68 ± 45**	75.67 ± 34.74**	58.17 ± 12.97**
Vitamin E (mg/dl)	1.09 ± 0.48	0.63 ± 0.37	1.15 ± 0.57	0.69 ± 0.10
GSH (mg/dl)	80.4 ± 9.4	78.3 ± 4.1	81.40 ± 12.1	74.2 ± 14.8
MDA (nm/ml)	3.7 ± 1.0	7.5 ± 1.4*	5.4 ± 1.0	5.5 ± 1.6
CD <sub>4</sub> /CD <sub>8</sub>	2.28 ± 1.10	1.30 ± 0.52*	2.24 ± 0.63	1.91 ± 0.83

\*\*p<0.01

\*p<0.05

RESULTS AND DISCUSSION

The values in the Table show that the drug abusers in general have considerable lower folate and beta carotene levels. The vitamin C status in both methadone and opiate groups were significantly lower than the control while the cocaine group showed a value that falls in the lower side of the normal range. Although

the vitamin E status was not statistically different, the values in the methadone and cocaine groups were borderline normal.

In summary, the methadone group has more nutrient shortages than either the heroin or cocaine abusers, showing low values for folate, and the antioxidant nutrients B-Carotene, vitamin C and vitamin E. Interestingly, the methadone group is the only one that showed significantly elevated levels of MDA indicating an abnormally high level of lipid peroxidation. This supports the notion that methadone treatment may be associated with an increased utilization of the body's antioxidant systems.

The finding that only the methadone-treated subjects have a lower lymphocyte CD<sub>4</sub>/CD<sub>8</sub> ratio (see Table) may suggest a relationship, direct or indirect, between their level of free radical production (i.e., as evidenced by a high lipid peroxidation and low levels of antioxidant nutrients) and their immune function. It is not clear whether the methadone treatment is the cause of the depletion of the antioxidant nutrients and immunodepression or whether the latter may result from the life-style and poor diet of these subjects.

Although the sample size in this preliminary study is small, the results presented here provide some significant information on the nutritional status of drug abusers. Since intravenous drug abusers have a higher risk for contracting HIV infection and since free radicals may be involved in the pathogenesis of viral infections, particularly AIDS (Papadopoulos-Eleopoulos, 1988), the potential interaction between nutritional status and

immunodepression in these subject becomes even more important. Cellular oxidative injury, measured as an increased production of malondialdehyde (Sonnerborg, et al., 1988) may play a role in the progression of HIV infection. A double-blind, placebo-controlled study with sodium diethyldithiocarbonate, a potent antioxidant, in patients with AIDS or AIDS-related complex have shown significant reduction in primary opportunistic infections (Lang, et al., 1988). Based on this information, a pertinent question may be asked: Will the nutritional and immunological status of HIV-infected drug abusers who enter into methadone maintenance programs deteriorate faster, thus leading to an earlier onset of full-blown AIDS disease? A larger, more thorough investigation of this relationship is, therefore, warranted.

#### REFERENCES

- BEUTLER E, DURON O, AND KELLY, BM. Improved Method for the Determination of Blood Glutathione. J. Lab. Clin. Med., 61:882-888, 1963.
- CASEY J, FLINN WR, YAO JST, FAHEY V, PAWLOWSKI J and BROGAN JJ. Correlation of Immune and Nutritional Status with Wound Complications in Patients Undergoing Vascular Operations. Surgery, 93:822-827, 1983.
- CHANDRA, RK. Nutrition, Immunity and Infection: Present Knowledge and Future Direction. Lancet, 1:688-691, 1983.
- DONAHOE RM and FALEK A. Neuroimmunomodulation by Opiates and Other Drugs of Abuse: Relationship to HIV infection and AIDS. In: Psychological, Neuropsychiatric and Substance Abuse Aspects of AIDS (eds. Bridge TP et al.) Raven Press, N.Y. 1988, pp. 145-152.
- FAHEY JL, PRINCE H, WEAVER M, GROOPMAN J, VISSCHER B, and SWARTZ RD. Quantitative Changes in T-Helper or T-Suppressor/Cytotoxic Lymphocyte Subsets that Distinguish Acquired Immune Deficiency Syndrom from other Immune Subset Disorders. Am. J. Med., 76:95-100, 1984.
- HERBERT JR, NICHOLS SE and KABAT GC. Indicators of Nutritional Status Among Clients from a New York City Methadone Treatment Center. J. Substance Abuse Treatment, 7:161-165, 1990.

KREEK MJ. Immune Function in Heroin Addicts and Former Heroin Addicts in Treatment: Pre/Post AIDS epidemic. In: Current Chemical and Pharmacological Advance on Drugs of Abuse Which Alter Immune function and Their Impact Upon HIV Infection. Pham. P.T.K. (ed.) NIDA Research Monograph Series. Rockville, MD, 96:192-219, 1990.

LANG JM, TOURAINE JL, TREPO C, CHOUTET P, KIRSTETTER M, FALKENRODT A, HERVIOU L, LIVROZET JM, RETORNAZ G, TOURAINE F, RENOUX G, and RENOUX M. Randomised Double-Blind Placebo-Controlled Trial of Ditiocarb Sodium (Imuthiol) in Human Immunodeficiency Virus Infection. Lancet ii:702-706, 1988.

MOHS ME, WATSON RR and LEONARD-GREEN T. Nutritional Effects of Marijuana, Heroin, Cocaine and Nicotine. J. Am. Diet. Assoc., 90:1261-1267, 1990.

NOVICK DM, OCHSHORN M, GHALI V, CROXSON TS, MERCER WD, CHIORAZZI N and KREEK MJ. Natural Killer Cell Activity and Lymphocyte Subsets in Parenteral Heroin Abusers and Long-term Methadone Maintenance Patients. J. Pharm. Exper. Ther., 250:606-610, 1989.

PAPADOPOULOS-ELEOPULOS, E. Reappraisal of AIDS: Is the Oxidation Induced by the Risk Factors the Primary Cause? Med. Hypoth., 25: 151-162, 1988.

SATOH K. Serum Lipid Peroxide in Cerebrovascular Disorders Determined by a New Calorimetric Method. Clinica Chimica Acta, 90:37-43, 1978.

SONNERBORG A, CARLIN G, AKERLUND B and JARSTRAND C. Increased Production of Malondialdehyde in Patients with HIV Infection. Scand. J. Infect. Dis., 20:287-290, 1988.

COPYRIGHT © 1991 BY

PJD Publications Ltd., P. O. Box 966, Westbury, NY 11590, U.S.A.

